

Primary care pathway: Peripheral arthritis

Quick links:

[Expanded details](#)

[Provider resources](#)

[Patient resources](#)

Peripheral joint pain

Red flags


Infection – fever, history of immunosuppression/IVDU, prosthetic joint
 Fracture – history of osteoporosis or risk factors, history of fall
 Malignancy – unexplained weight loss, significant night pain, severe fatigue, history of cancer

Yes

Refer to ER/page rheumatologist on call if red flags present or if your patient needs to be seen immediately

No

Where is your patient's joint pain?
(refer to diagrams)



Probable osteoarthritis
If pain pattern matches the joints highlighted in **left** diagram, patient likely has osteoarthritis

Probable inflammatory arthritis
If pain pattern matches the joints highlighted in **right** diagram, screen for inflammatory arthritis

Are there typical features of osteoarthritis?

- Morning stiffness lasting <30 mins
- Symptoms worsening with use
- Symptoms worsening throughout the day
- Typical joints: carpometacarpal (CMC), knee, hip
- Gradual bony joint enlargement

No

Proceed with work-up of inflammatory arthritis

Yes

CCFP/AAC toolkit
If conservative management fails, consider these investigations:

- CBC, CRP, TSH, Albumin, Calcium, +/- urate
- X-rays of affected joint

Does your patient have inflammatory features?

- Morning stiffness lasting >30 mins
- Acute/New onset joint swelling/effusion
- 3 or more joints
- Symptoms improving with activity/worsening with rest
- Symptoms improving throughout day (may worsen at end of day)

Does your patient have extra-articular features?

- Psoriasis
- Inflammatory bowel disease
- Uveitis/iritis
- Urethritis, diarrheal illness
- Rash (butterfly/photosensitivity)
- Raynaud's
- Nodules
- Family history of autoimmune disorder

Investigations

- CBC, Cr, UA, TSH, calcium, albumin, CRP, +/- urate, RF, anti-CCP, ANA
- X-ray symptomatic joints +/- x-ray lumbar/spine and SI joints

Does your patient have inflammatory back pain?

- Morning back stiffness >30 mins
- Improves w/activity, worsens w/rest
- Alternating buttock pain
- Nocturnal waking due to pain in second half of night
- Improves with NSAIDs

If "yes" for any of the above questions, your patient may have inflammatory arthritis. Refer to rheumatology and/or call Specialist Link.

[Rheumatology access](#)

[Specialist Link](#)

EXPANDED DETAILS

Peripheral arthritis is grouped into non-inflammatory and inflammatory arthritides. Non-inflammatory arthritis, most commonly osteoarthritis, is the most frequent form of peripheral arthritis. This form of arthritis can be safely managed in the medical home. Inflammatory arthritis is less common, occurring in approximately 1/100 patients. This form of arthritis, when active, requires a semi-urgent assessment by a rheumatologist (within approximately 6-8 weeks).

Osteoarthritis can affect any joint, but most commonly affects the following joints: the hands (nodal osteoarthritis), carpal metacarpal (CMC), knees, and hips. Risk factors for this form of arthritis include age > 40, female gender, increased BMI, family history and prior joint injury. This type of arthritis is characterized by pain with <30 minutes of am stiffness. Symptoms tend to worsen throughout the day and with activity. This form of arthritis can usually be managed in the medical home with lifestyle modifications (weight loss), pain control (acetaminophen, NSAIDs, intra-articular steroid or hyaluronic acid injections through interventional radiology), physiotherapy and occupational interventions (use of a cane, off-loading knee brace). Management of these patients may be assisted with the use of the CCFP toolkit at [Osteoarthritis tool \(cfpc.ca\)](#). Imaging may be helpful in evaluating patients with mechanical joint pain. (see section: *Radiographs recommended for work-up (if needed) of peripheral arthritis*) X-ray findings show joint space narrowing, osteophytes, subchondral sclerosis/cysts. Periarticular osteopenia is absent.

Erosive osteoarthritis is a rare form of osteoarthritis in which patients' osteoarthritis may present with inflammatory arthritis symptoms. X-ray findings may reveal central erosions. Patients with erosive osteoarthritis may benefit from a rheumatology consultation.

Inflammatory arthritis is an autoimmune condition that, if left untreated, can lead to significant joint damage and disability. Risk factors for inflammatory arthritis include a family history of inflammatory arthritis, first nations ethnicity, a history of psoriasis, new or severe Raynaud's, rash or recent urethritis/diarrheal illness, a personal or family history of spondyloarthropathy features (psoriasis, uveitis/iritis, ankylosing spondylitis, Crohn's disease or ulcerative colitis, reactive arthritis). This form of arthritis is characterized by joint pain and *swelling with >30 minutes of morning stiffness*. Patients often describe much longer morning stiffness, sometimes lasting most of the day. Symptoms *improve throughout the day and with activity*. Patients may have a single joint (monoarthritis), a few joints (oligoarthritis) or many joints (polyarthritis) involved. Joints involved may be large (knee, ankle, shoulder) or small (wrist, MCPs, PIPs, MTPs). Patients may have nodules or joint deformities in the advanced stages of the disease. Inflammatory arthritides include seropositive and seronegative rheumatoid arthritis (RA), psoriatic arthritis, ankylosing spondylitis, inflammatory bowel disease-related arthritis, systemic lupus erythematosus and scleroderma.

When an inflammatory arthritis is suspected, further investigations should be ordered. Some of these include CBC, creatinine, urinalysis, urate, rheumatoid factor (RF), anti-citrullinated c-peptide (anti-CCP) and c-reactive protein (CRP) (see section: *Common labs useful in the work-up of peripheral inflammatory arthritis*). If there are specific concerns regarding a connective tissue disease or Lupus, an ANA may be helpful. It is important to note that an ANA of 1:80 is considered "negative" no matter what the pattern. Patients with inflammatory arthritis commonly have a normocytic anemia, thrombocytosis, and elevated inflammatory markers. RA patients may have a positive RF or anti-CCP. X-rays of the affected joints should be ordered (see section: *Radiographs recommended for work-up (if needed) of peripheral arthritis*) with consideration of the lumbar spine and bilateral sacroiliac joints ONLY if co-morbid



inflammatory back pain is suspected (i.e. Ankylosing spondylitis). X-rays may show periarticular osteopenia, periostitis, or erosive changes; however, they may also be normal, particularly in early disease. When a patient presents with symptoms that suggest an inflammatory arthritis, he or she should be referred semi-urgently to a rheumatologist for further work-up and management as early intervention can alter the natural course of the disease.

Occasionally, patients with an inflammatory arthritis may present with a monoarthritis. In these cases, a diagnosis of gout or septic arthritis must be considered. Red flags for septic arthritis include rapid onset and systemic symptoms such as fevers. Risk factors for septic arthritis include age > 80, diabetes mellitus or co-existent rheumatoid arthritis, prosthetic joint or recent joint surgery and skin infection. Septic arthritis is a medical emergency and, when it is suspected, patients should be sent directly to the nearest emergency department. Risk factors for gout include a history of podagra, family history of gout, chronic kidney disease, hypertension, and nephrolithiasis, the presence of hyperuricemia, and the presence of gouty tophi on physical examination. Gout may also be safely managed in the medical home. *For further information on the work-up and management of gout, please see the enhanced primary care pathway for gout at specialistlink.ca.*

Checklist to guide your in-clinic review of this patient with peripheral joint pain:

| |
|---|
| <input type="checkbox"/> Absence of red flags |
| <input type="checkbox"/> Use of the clinical flow diagram to differentiate whether patient’s presentation is consistent with peripheral inflammatory or non-inflammatory arthritis. |
| <input type="checkbox"/> Patient’s presentation is consistent with peripheral <i>non-inflammatory arthritis</i> : see the focused summary in this pathway or the CCFP Toolkit at Osteoarthritis tool (cfpc.ca) for management. |
| <input type="checkbox"/> Patient presentation is consistent with peripheral <i>non-inflammatory arthritis, but has failed to improve with the interventions suggested by the CCFP Toolkit?</i> Consider investigations as outlined in Section 5 and 6; and/or calling Specialist Link for further advice. |
| <input type="checkbox"/> Patient’s presentation is consistent with peripheral <i>inflammatory arthritis</i> : consider investigations as outlined in section 5 and 6. |
| <input type="checkbox"/> Patient’s presentation and investigations support the diagnosis of <i>inflammatory arthritis</i> : use access pathway to refer semi-urgently to Rheumatology as early intervention can alter the course of the disease. |



Common labs useful in the work-up up peripheral inflammatory arthritis

Complete blood count (CBC) – Anemia and thrombocytosis may be subtle signs of inflammation.

C-reactive protein (CRP) – Often elevated in the setting of inflammatory disorders.

Creatinine (Cr) – Some disorders such as connective tissue diseases, may have renal involvement. Creatinine is also useful in the dosing of some medications used in the management of arthritis.

Urinalysis (UA) – Some inflammatory arthritides are associated with renal dysfunction. The presence of proteinuria and hematuria, will trigger the work-up systemic disorders such as lupus and vasculitis.

Urate – Elevated uric acid is often, but not always, associated with gout. Many patients with hyperuricemia do not ever develop gout, therefore, it is not diagnostic of gout. It should be noted that during an acute gout attack, the urate may be low or normal.

Thyroid stimulating hormone (TSH) – Joint pain may be a manifestation of hypothyroidism and resolves with treatment of the thyroid disorder. Hypothyroidism is also a risk factor for pseudo gout.

Corrected Calcium (Calcium/albumin) – Hypercalcemia is often associated with joint pain. Recall “bones, groans, psychological overtones.” In addition, hypercalcemia may put patients at risk of pseudo gout.

Rheumatoid Factor (RF) – Rheumatoid Factor (RF) is commonly associated with rheumatoid arthritis; however, it has a sensitivity of only 66% and a specificity of only 82% for rheumatoid arthritis. It is also associated with other systemic diseases such as Sjogren’s Syndrome and cryoglobulinemic vasculitis. It can also be present in normal healthy individuals. A higher titre is more predictive of an underlying disease process. RF can be present in patients up to 10 years prior to onset of symptoms.

Anti-citrullinated c-peptide (anti-CCP/ACPA) – This antibody is more sensitive and specific for rheumatoid arthritis, than RF. Specifically, the sensitivity of anti-CCP is 70% and the specificity is 95%. It predicts more aggressive and erosive disease and is useful to prognosticate and triage patients. Anti-CCP/ACPA may be present up to 10 years prior to onset of disease symptoms. A positive anti-CCP/ACPA indicates significantly increases the pre-test probability of rheumatoid arthritis. Patients with a positive anti-CCP/ACPA should be referred to rheumatology.

Anti-nuclear antibody (ANA) – A positive ANA is associated with a variety of connective tissue diseases including lupus, Sjogren’s, myositis and scleroderma. Both the titre and the pattern of the ANA may be useful in elucidating the underlying cause of the arthritis. It is important to note that an ANA titre of 1:80 is considered negative, no matter what the pattern. In addition, approximately 5% of normal healthy adults (a higher percentage in pediatric patients) have a positive ANA. While useful in the work up, a positive ANA is not diagnostic of an autoimmune disease. However, in the presence of peripheral joint pain, it may require further investigation. ANA is positive in up to 20-30% of RA patients.



Radiographs recommended (if needed) for work-up of peripheral arthritis

Radiographs may not be necessary in the initial work up of both inflammatory and osteo-arthritis. Scenarios in which they may be considered include: the need to confirm the diagnosis in uncertain cases; cases in which the patient has failed to respond to evidence-based treatment (i.e. CCFP Toolkit for Management of OA); staging disease severity: or when considering a rheumatology or orthopedic surgery consultation. **Please be specific when ordering views.** The following radiographs are recommended for the initial assessment of peripheral arthritis (*non-traumatic* joint pain) if warranted.

SHOULDER

- AP (anterior-posterior), glenoid, axillary

ELBOW

- AP, lateral, both obliques

WRIST

- AP, lateral, oblique

HAND/FINGERS

- AP, lateral, oblique

HIP

- Weightbearing AP pelvis, AP and lateral of affected hip

KNEE

- Bilateral weightbearing AP and tunnel views (PA flexed 30 degrees), lateral and skyline of affected joint

ANKLE

- AP, mortise, lateral

FOOT/TOES

- AP, lateral, oblique

*Additional special x-ray views can be ordered with appropriate supportive clinical history.

****MRI is NOT indicated for chronic joint pain and x-ray confirmed osteoarthritis***

*Ultrasound has a very limited role in chronic knee pain, and is generally used to evaluate for: joint effusion pre-arthrocentesis, integrity of the extensor mechanism (quads and patellar tendons), and solid vs. cystic masses (including Baker's cyst)

****Ultrasound is NOT indicated for the routine evaluation of osteoarthritis.***

****Ultrasound is NOT indicated for assessing menisci or cruciate ligaments of the knee.***



BACKGROUND

About this pathway

- The pathway is intended to provide evidence-based guidance to support primary care and specialty care providers in caring for adult patients with Peripheral arthritis within the medical home. Rheumatology tele-advice is available via [Specialist Link](#) (see advice section under Provider Resources).

Authors and conflict of interest declaration

- This pathway was developed by leveraging the collective knowledge, experience and expertise of many individuals. See a full list below. For more information, please email info@calgaryareapcns.ca.

| Faculty | Names |
|--|--------------------------------|
| Professor of Medicine, Division Chief Section of Rheumatology University of Calgary | Dianne Mosher, MD, FRCPC |
| Clinical Assistant Professor Section of Rheumatology University of Calgary | Corisande Baldwin, MD, FRCPC |
| Clinical Assistant Professor Section Chief Community Primary Care Department of Family Medicine-Calgary Zone | Monica Sargious MD, CCFP, FCFP |
| Clinical Associate Professor Department of Radiology University of Calgary | Bevan Frizzell, MD, FRCPC |
| Medical Director, Calgary West Central - PCN | Jane Ballantine, MD, CCFP |
| Clinical Assistant Professor Section of Rheumatology University of Calgary | Maysan Abu-Hakima, MD, FRCPC |
| Medical Director, Mosaic Primary Care Network | Oliver David MD, CCFP |

Pathway review process, timelines

- Primary care pathways undergo scheduled review every three years, or earlier if there is a clinically significant change in knowledge or practice. This pathway was developed in May 2018 and was most recently updated in May 2024. The next scheduled review is May, 2026. However, we welcome feedback at any time. Please email comments to info@calgaryareapcns.ca with "Peripheral arthritis Pathway" in the subject line.

Copyright information

This work is licensed under a Creative Commons Attribution-Non-Commercial-Share Alike 4.0 International license. You are free to copy, distribute and adapt the work for non-commercial purposes, as long as you attribute the work to Alberta Health Services and Primary Care Networks and abide by the other license terms. If you alter, transform, or build upon this work, you may distribute the resulting work only under the same, similar, or compatible license. The license does not apply to content for which the Alberta Health Services is not the copyright owner.



DISCLAIMER

This pathway represents evidence-based best practice but does not override the individual responsibility of health care professionals to make decisions appropriate to their patients using their own clinical judgment given their patients' specific clinical conditions, in consultation with patients/alternate decision makers. The pathway is not a substitute for clinical judgment or advice of a qualified health care professional. It is expected that all users will seek advice of other appropriately qualified and regulated health care providers with any issues transcending their specific knowledge, scope of regulated practice or professional competence.



PROVIDER RESOURCES

Advice options

- Non-urgent telephone advice connects family physicians, nurse practitioners and specialists in real time via a tele-advice line. Family physicians, nurse practitioners and specialists can request non-urgent advice from a rheumatologist, at [specialistlink.ca](http://www.specialistlink.ca) or by calling 403-910-2551. This service is available from 8 a.m. to 5 p.m. Monday to Friday (excluding statutory holidays). Calls are returned within one (1) hour.
- Non-urgent rheumatology advice is available across the province via Alberta Netcare eReferral Advice Request (responses are received within five calendar days). View <https://www.albertanetcare.ca/eReferral.htm> for more information.

| Resources | Location |
|---|---|
| Clinical Pathways | http://www.specialistlink.ca/ |
| RheumInfo — Accurate information on arthritis and its medications | https://rheuminfo.com/ |
| ArthritisID PRO app | http://www.arthritisresearch.ca/tools/apps |
| Arthritis Research Canada: Osteoarthritis Knee Examination | https://www.youtube.com/watch?v=BS8KbciQSuk |
| CCFP Toolkit for Management of Osteoarthritis | Osteoarthritis tool (cfpc.ca) |

PATIENT RESOURCES

Information

| Resource type | Resource name | Location |
|---------------|---|---|
| Website | RheumInfo — Accurate information on arthritis and its medications | https://rheuminfo.com/ |
| Website | The Arthritis Society of Canada | http://arthritis.ca/ |
| Website | Arthritis ID app | http://www.arthritisresearch.ca/tools/apps |

